



## P.44 Artificial DNA membrane nanopores

K Göpfrich<sup>1</sup>, T Zettl<sup>1</sup>, J R Burns<sup>2</sup>, V V Thacker<sup>3</sup>, S Hernández-Ainsa<sup>1</sup>, N A W Bell<sup>1</sup>, E Stulz<sup>3</sup>, S Howorka<sup>2</sup> and U F Keyser<sup>1</sup>

<sup>1</sup>Cavendish Laboratory, University of Cambridge, UK, <sup>2</sup>Department of Chemistry, University College London, UK,

<sup>3</sup>School of Chemistry, University of Southampton, UK

Membrane nanopores are essential components of biological and artificial cells. Our group has shown that we can create artificial nanopores using DNA origami self-assembly [1], [2] and anchor them in lipid membranes [3]. Insertion of negatively charged DNA pores into a hydrophobic membrane is achieved by attaching functional hydrophobic groups in strategic positions on the DNA nanopores. Pore formation in lipid vesicles is studied for different nanopore designs and hydrophobic modifications via fluorescent imaging [4]. We demonstrate membrane anchoring using only two porphyrin-based lipid anchors. The porphyrin moieties also act as fluorescent dyes to aid the microscopic visualization of the DNA nanopore [3]. Single-channel current recordings of our artificial DNA nanopores are performed using a high-throughput lipid nanobilayer system that has recently been introduced by our group [5], [6]. Pore architecture and functionality of our DNA nanopores can be easily adapted, opening the pathway to design novel membrane channels.

- [1] Bell, N. a W., Engst, C. R., Ablay, M., Divitini, G., Ducati, C., Liedl, T., & Keyser, U. F. (2012). DNA origami nanopores. *Nano Letters*, 12(1), 5127
- [2] Hernández-Ainsa, S., Bell, N. A. W., Thacker, V. V, Göpfrich, K., Misiunas, K., Fuentes-Perez, M. E., Moreno Herrero, F. & Keyser, U. F.(2013). DNA origami nanopores for controlling DNA translocation. *ACS Nano*, doi:10.1021/nn401759r
- [3] Burns, J. R., Göpfrich, K., Wood, J. W., Thacker, V. V., Stulz, E., Keyser, U. F., & Howorka, S. (2013). Lipid Bilayer-Spanning DNA Nanopores with a Bifunctional Porphyrin Anchor. *Angewandte Chemie*. Published as a 'Hot Paper', Front cover article
- [4] Thacker, V. V, Ghosal, S., Hernández-Ainsa, S., Bell, N. a W., & Keyser, U. F. (2012). Studying DNA translocation in nanocapillaries using single molecule uorescence. *Applied Physics Letters*, 101(22), 223704
- [5] Göpfrich, K., Kulkarni, C. V, Pambos, O. J., & Keyser, U. F. (2013). Lipid Nanobilayers to Host Biological Nanopores for DNA Translocations. *Langmuir*, 29(1), 355364
- [6] Gornall, J. L., Mahendran, K. R., Pambos, O. J., Steinbock, L. J., Otto, O., Chimere, C., Winterhalter, M. & Keyser, U. F. . (2011). Simple reconstitution of protein pores in nano lipid bilayers. *Nano Letters*, 11(8), 333440