



Poster session 2 – Tuesday 5 July

P2.068 IBD background rejection and tagging at the Double Chooz experiment

A Meregaglia¹ and A Hourlier²

¹CNRS / IPHC, France, ²Laboratoire AstroParticules et Cosmologie Université Paris Diderot, France

on behalf of Double Chooz collaboration

Double Chooz (DC) is a reactor anti-neutrino disappearance experiment located in France near the power plant of Chooz. The main goal of the experiment is the measurement of the θ_{13} mixing angle and, already in 2011, it provided the first positive indication for a non-zero value of such an oscillation parameter, using a single far detector. The exploitation of the near detector, taking data since December 2014, provided the partial cancellation of some systematic errors. We show the first results for both detectors, including both phases: single detector (2011-2014) and double detector (from 2015). Major improvements have followed through the different released analyses, especially on the reduction and active rejection of both radiogenic and cosmogenic backgrounds. Currently, DC has among the best-correlated and uncorrelated (accidental) background systematics errors in the θ_{13} reactor field. In fact, DC is able to actively reject all types of backgrounds using several unprecedented multidetector correlation strategies. So, multi-Compton radioactive, stopping muon, fast-neutron and even the generation of spallation isotopes such as Li(9), H(8) and B(12) can be tagged, for energy spectra characterisation and eventual rejection. In addition, the FADC based electronics provide the recording of the waveform of each PMT, which is exploited for additional background rejection based on pulse shape analyses in time and frequency domain (i.e. Fourier transformed). All the strategies here described are validated by invaluable reactor-OFF data, only available to DC (1 week). Last, upon the measurement of θ_{13} , DC articulates further constraining of background exploiting its excellent response control, reducing the main background errors by up to 2x. In this poster, we illustrate the full background tagging and rejection framework, including the articulation of background higher precision improvement during neutrino oscillation fit. The hereby presented techniques are expected to be of large value for the development of similar techniques by similar detectors.