



Poster session 4 – Friday 8 July

P4.074 Search for double beta decay processes of ^{124}Xe with XENON100 & XENON1T

A Fieguth

WWU Muenster, Germany

on behalf of XENON collaboration

Driven by the search for dark matter particles the XENON project installed its next stage multi-ton experiment XENON1T at LNGS, which will probe the spin-independent-WIMP nucleon cross section down to $2 \times 10^{-47} \text{cm}^2$. Besides its main purpose different particle physics topics can be addressed. One example are double beta decay processes of the natural isotope ^{124}Xe . It is expected to decay via two-neutrino double electron capture (2νECEC). Due to its high Q-value of 2864 keV the positron including processes $2\nu\text{EC}\beta^+$ and $2\nu\beta^+\beta^+$ are also allowed. In absence of a detection there is only a lower limit to the respective half-life (e.g. $> 4.7 \times 10^{21} \text{yr}$ for 2νECEC). With its fiducial mass of one ton XENON1T will be the most sensitive detector for these decays to date and a detection the 2νECEC and $2\nu\text{EC}\beta^+$ decays is within reach providing information on nuclear matrix element (NME) models. In addition to these standard model processes the detection of a neutrinoless decay (e.g. 0νECEC) would hint towards physics beyond the Standard Model and could shed light on the neutrino mass and a Majorana nature of neutrinos. Although a resonance enhancement is possible for this particular decay, the half-life predictions are ranging from 10^{24}yr to 10^{31}yr illustrating the high uncertainty for calculations of the NME involved in this decay as well as the unknown neutrino mass value. Due to emission of an internal bremsstrahlung photon accompanied by secondary particles (e.g. Auger electrons) a coincidence signature provides a background suppression to the already low background level of the XENON1T experiment and, therefore, the lower bound of the predictions could be within reach. Until XENON1T data are available, the search for all processes can be performed in the data of its predecessor XENON100, which could compete with existing searches. This work is supported by BMBF under contract number 05A14PM1 and DFG (GRK 2149).